

(iii) 0.6

(iv) 0.8

Q.2 Attempt Any Two

[10]

(a) If A and B are two events such that $P(A) = \frac{3}{8}$, $P(B) = \frac{5}{8}$ and $P(A \cup B) = \frac{3}{4}$,

5 A 4

find $P(A/B)$ and $P(B/A)$. Check whether A and B are independent.

(b) In a bolt factory, machines A , B and C manufacture 25%, 35% 40% of the total output and out of the total manufacturing, 5%, 4% and 2% bolts are defective. A bolt is drawn at random from the product and it is found to be defective. Find the probability that it is manufactured by machine A .

5 A 4

(c) A discrete random variable has the probability mass function given below:

5 A 4

X	-2	-1	0	1	2	3
$P(X = x)$	0.2	k	0.1	$2k$	0.1	$2k$

Find k , mean and variance.

Q.3 Attempt Any Two

[10]

(a) Fit a curve $y = a + bx$ to the following data:

5 A 3

x	0	1	2	3	4
y	1	1.8	3.3	4.5	6.3

(b) Fit a least square second degree quadratic curve $y = a + bx + cx^2$ to the following data:

5 A 3

x	1	2	3	4
y	1.7	1.8	2.3	3.2

(c) Fit a curve $y = ab^x$ to the following data:

5 A 3

x	1	2	3	4	5	6	7	8
y	1	1.2	1.8	2.5	3.6	4.7	6.6	9.1

Q.4 Attempt Any Two

[10]

(a) Calculate the first four moments from the following data:

5 A 5

x	0	1	2	3	4	5	6	7	8
f	5	10	15	20	25	20	15	10	5

Also, calculate the values of β_1 and β_2 .

(b) Calculate the standard deviation of the following data:

5 A 5

x	10	11	12	13	14	15	16	17	18
f	2	7	10	12	15	11	10	6	3

(c) The runs scored by two batsmen A and B in 9 consecutive innings are given below:

5 A 5

A	85	20	62	28	74	5	69	4	13
B	72	4	15	30	59	15	49	27	26

Which batsman is more consistent?

SECTION B

Marks BL CO

Q.5 Multiple-Choice Questions

[05]

- (a) Interpolation is a method of _____. 1 R 2
- (i) Interrelating (ii) Estimating
- (iii) Integrating (iv) combining
- (b) Which method can be used for both equal and unequal intervals? 1 R 2
- (i) Newton's method (ii) Cubic spline method
- (iii) Lagrange's Method (iv) None of the above
- (c) $\nabla \log x =$ _____. 1 R 2
- (i) $\log \frac{x}{x-h}$ (ii) $\log \frac{x-h}{x}$
- (iii) $\log \frac{x}{x+h}$ (iv) $\log(x-h)$
- (d) The complete solution of $z = px + qy + pq$ is _____. 1 R 4
- (i) $z = ax + by$ (ii) $z = ax + by + ab$
- (iii) $z = ax + by + \log(ab)$ (iv) none of these
- (e) What is the nature of Lagrange's linear partial differential equation? 1 R 4
- (i) First order, Third degree (ii) Second order, First degree
- (iii) First order, Second degree (iv) First order, First degree

Q.6 Attempt Any Two

[10]

- (a) Use Newton's forward difference method to find the approximate value of $f(2.3)$ from the following data: 5 A 2

x	2	4	6	8
$f(x)$	4.2	8.2	$\frac{12.}{2}$	16.2

- (b) The population of a town is given below. Estimate the population for the year 1930 using suitable interpolation. 5 A 2

Year	x	1891	1901	1911	1921	1931
Population	y	46	66	81	93	101
(in thousand)						

(c) By using Lagrange's formula, find y when $x = 10$. 5 A 2

x	5	6	9	11
y	12	13	14	16

Q.7 Attempt Any Two [10]

(a) Using Newton's divided difference interpolation, compute the value of $f(6)$ from the table given below: 5 A 2

x	1	2	7	8
y	1	5	5	4

(b) Construct Newton's forward interpolation polynomial for the following data: 5 A 2

x	4	6	8	10
y	1	3	6	16

(c) Use Lagrange's formula to fit a polynomial to the data: 5 A 2

x	-1	0	2	3
y	8	3	1	12

and hence, find $y(2)$.

Q.8 Attempt Any Two [10]

(a) Eliminate the function f from the relation $f(xy + z^2, x + y + z) = 0$. 5 A 1

(b) Solve $yzp - xzq = xy$. 5 A 1

(c) Solve (i) $p + q^2 = 1$ (ii) $p^2 - q^2 = x - y$. 5 A 1
